

Computational Linguistics: Semantics

Raffaella Bernardi

14-10-2010

1 Exercise 1a: From Relational to Functional Perspective

Look at the Knowledge Base below.

1. Harry is a wizard.
2. Hagrid scares Dudley.
3. All wizards are magical.
4. Uncle Vernon hates anyone who is magical.
5. Aunt Petunia hates anyone who is magical and scares Dudley.

Build a model for it by (i) writing your interpretation for *wizards*, *magical*, *scares*, *hates* using the relational interpretation first, and then the functional one.

Solution:

$\llbracket \textit{wizard} \rrbracket$	$=$	$\{\textit{harry}\}$	$\{x \mid \textit{wizard}(x) = 1\}$
$\llbracket \textit{magical} \rrbracket$	$=$	$\{\textit{harry}\}$	$\{x \mid \textit{magical}(x) = 1\}$
$\llbracket \textit{scares} \rrbracket$	$=$	$\{(\textit{hagrid}, \textit{dudley})\}$	$\{(x, y) \mid (\textit{scares } (y)) (x)\}$
$\llbracket \textit{hates} \rrbracket$	$=$	$\{(\textit{vernon}, \textit{harry})\}$	$\{(x, y) \mid (\textit{hates } (y)) (x)\}$

(ii) Specifying the types of the expressions in your universe, and (iii) the domains of interpretation. E.g.

The domain of entities is as below:

$$D_e = \{\textit{harry}, \textit{hagrid}, \textit{vernon}, \textit{petunia}, \textit{dudley}\}$$

Solution: $D_{(e \rightarrow t)} = \{\textit{magical}, \textit{wizard}\}$, $D_{(e \rightarrow (e \rightarrow t))} = \{\textit{scare}, \textit{hates}\}$

2 Exercise 2: Well formed formula

Let j be a constant of type e ; M of type $e \rightarrow t$; S of type $((e \rightarrow t) \rightarrow (e \rightarrow t))$, and P of type $(e \rightarrow t) \rightarrow t$. Furthermore, x is a variable of type e , and Y a variable of type $(e \rightarrow t)$.

Determine which of the following is well-formed, give its type.

1. $(\lambda x.M(x))(P)$. [NWF]

2. $(\lambda x.M(x))(j)$. [WF]
3. $\lambda x.M(j)$. [NWF: vacus abstraction]
4. $S(\lambda x.M(x))$. [WF]
5. $(\lambda Y.Y(j))(M)$ [WF]
6. $\lambda x.(M(x) \wedge M(j))$ [WF]
7. $(\lambda x.M(x)) \wedge M(j)$ [NWF: $\lambda x.M(x)$ and $M(j)$ are of types $e \rightarrow t$ and t , resp. \wedge coordinates terms of types t]

3 Exercise 3: β -conversion

Let j be a constant of type e ; M of type $(e \rightarrow t)$, and A of type $e \rightarrow (e \rightarrow t)$. Furthermore, x and y are variables of type e , and Y is a variable of type $e \rightarrow t$. Reduce the following expression as much as possible by means of β -conversion.

1. $\lambda x(M(x))(j)$ [M(j)]
2. $\lambda Y(Y(j))(M)$ [M(j)]
3. $\lambda x\lambda Y(Y(x))(j)(M)$ [M(j)]
4. $\lambda x\forall y(A(x)(y))(j)$ [$\forall y.A(j)(y)$]
5. $\lambda x\forall y(A(x)(y))(y)$ [$\forall y.A(z)(y)$]
6. $\lambda Y(Y(j))\lambda x(M(x))$ [M(j)]
7. $\lambda Y\forall x(Y(x))\lambda y(A(x)(y))$ [$\forall z.A(x)(z)$]