

## Discrete random variables

### Probability mass function

Given a discrete random variable  $X$  taking values in  $\mathcal{X} = \{v_1, \dots, v_m\}$ , its *probability mass function*  $P : \mathcal{X} \rightarrow [0, 1]$  is defined as:

$$P(v_i) = \Pr[X = v_i]$$

and satisfies the following conditions:

- $P(x) \geq 0$
- $\sum_{x \in \mathcal{X}} P(x) = 1$

## Discrete random variables

### Expected value

- The *expected value*, *mean* or *average* of a random variable  $x$  is:

$$E[x] = \mu = \sum_{x \in \mathcal{X}} xP(x) = \sum_{i=1}^m v_i P(v_i)$$

- The *expectation operator* is linear:

$$E[\lambda x + \lambda' y] = \lambda E[x] + \lambda' E[y]$$

### Variance

- The *variance* of a random variable is the moment of inertia of its probability mass function:

$$\text{Var}[x] = \sigma^2 = E[(x - \mu)^2] = \sum_{x \in \mathcal{X}} (x - \mu)^2 P(x)$$

- The *standard deviation*  $\sigma$  indicates the typical amount of deviation from the mean one should expect for a randomly drawn value for  $x$ .

## Properties of mean and variance

### second moment

$$E[x^2] = \sum_{x \in \mathcal{X}} x^2 P(x)$$

### variance in terms of expectation

$$\text{Var}[x] = E[x^2] - E[x]^2$$

### variance and scalar multiplication

$$\text{Var}[\lambda x] = \lambda^2 \text{Var}[x]$$

### variance of uncorrelated variables

$$\text{Var}[x + y] = \text{Var}[x] + \text{Var}[y]$$

## Probability distributions

### Bernoulli distribution

- Two possible values (outcomes): 1 (success), 0 (failure).
- Parameters:  $p$  probability of success.
- Probability mass function:

$$P(x; p) = \begin{cases} p & \text{if } x = 1 \\ 1 - p & \text{if } x = 0 \end{cases}$$

- $E[x] = p$
- $\text{Var}[x] = p(1 - p)$

*Example: tossing a coin*

- Head (success) and tail (failure) possible outcomes
- $p$  is probability of head

### Bernoulli distribution

#### Proof of mean

$$\begin{aligned} E[x] &= \sum_{x \in \mathcal{X}} xP(x) \\ &= \sum_{x \in \{0,1\}} xP(x) \\ &= 0 \cdot (1 - p) + 1 \cdot p = p \end{aligned}$$

### Bernoulli distribution

#### Proof of variance

$$\begin{aligned} \text{Var}[x] &= \sum_{x \in \mathcal{X}} (x - \mu)^2 P(x) \\ &= \sum_{x \in \{0,1\}} (x - p)^2 P(x) \\ &= (0 - p)^2 \cdot (1 - p) + (1 - p)^2 \cdot p \\ &= p^2 \cdot (1 - p) + (1 - p) \cdot (1 - p) \cdot p \\ &= (1 - p) \cdot (p^2 + p - p^2) \\ &= (1 - p) \cdot p \end{aligned}$$

## Probability distributions

### Binomial distribution

- Probability of a certain number of successes in  $n$  independent Bernoulli trials
- Parameters:  $p$  probability of success,  $n$  number of trials.
- Probability mass function:

$$P(x; p, n) = \binom{n}{x} p^x (1-p)^{n-x}$$

- $E[x] = np$
- $\text{Var}[x] = np(1-p)$

*Example: tossing a coin*

- $n$  number of coin tosses
- probability of obtaining  $x$  heads

### Pairs of discrete random variables

#### Probability mass function

Given a pair of discrete random variables  $X$  and  $Y$  taking values  $\mathcal{X} = \{v_1, \dots, v_m\}$   $\mathcal{Y} = \{w_1, \dots, w_n\}$ , the *joint probability mass function* is defined as:

$$P(v_i, w_j) = \Pr[X = v_i, Y = w_j]$$

with properties:

- $P(x, y) \geq 0$
- $\sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} P(x, y) = 1$

#### Properties

- Expected value

$$\mu_x = E[x] = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} x P(x, y)$$

$$\mu_y = E[y] = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} y P(x, y)$$

- Variance

$$\sigma_x^2 = \text{Var}[(x - \mu_x)^2] = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} (x - \mu_x)^2 P(x, y)$$

$$\sigma_y^2 = \text{Var}[(y - \mu_y)^2] = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} (y - \mu_y)^2 P(x, y)$$

- Covariance

$$\sigma_{xy} = E[(x - \mu_x)(y - \mu_y)] = \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} (x - \mu_x)(y - \mu_y) P(x, y)$$

- Correlation coefficient

$$\rho = \frac{\sigma_{xy}}{\sigma_x \sigma_y}$$

## Probability distributions

### Multinomial distribution (one sample)

- Models the probability of a certain outcome for an event with  $m$  possible outcomes.
- Parameters:  $p_1, \dots, p_m$  probability of each outcome
- Probability mass function:

$$P(x_1, \dots, x_m; p_1, \dots, p_m) = \prod_{i=1}^m p_i^{x_i}$$

- where  $x_1, \dots, x_m$  is a vector with  $x_i = 1$  for outcome  $i$  and  $x_j = 0$  for all  $j \neq i$ .
- $E[x_i] = p_i$
- $\text{Var}[x_i] = p_i(1 - p_i)$
- $\text{Cov}[x_i, x_j] = -p_i p_j$

## Probability distributions

### Multinomial distribution: example

- Tossing a dice with six faces:
  - $m$  is the number of faces
  - $p_i$  is probability of obtaining face  $i$

## Probability distributions

### Multinomial distribution (general case)

- Given  $n$  samples of an event with  $m$  possible outcomes, models the probability of a certain distribution of outcomes.
- Parameters:  $p_1, \dots, p_m$  probability of each outcome,  $n$  number of samples.
- Probability mass function (assumes  $\sum_{i=1}^m x_i = n$ ):

$$P(x_1, \dots, x_m; p_1, \dots, p_m, n) = \frac{n!}{\prod_{i=1}^m x_i!} \prod_{i=1}^m p_i^{x_i}$$

- $E[x_i] = np_i$
- $\text{Var}[x_i] = np_i(1 - p_i)$
- $\text{Cov}[x_i, x_j] = -np_i p_j$

## Probability distributions

*Multinomial distribution: example*

- Tossing a dice
  - $n$  number of times a dice is tossed
  - $x_i$  number of times face  $i$  is obtained
  - $p_i$  probability of obtaining face  $i$

## Conditional probabilities

**conditional probability** probability of  $x$  once  $y$  is observed

$$P(x|y) = \frac{P(x, y)}{P(y)}$$

**statistical independence** variables  $X$  and  $Y$  are statistical independent iff

$$P(x, y) = P(x)P(y)$$

implying:

$$P(x|y) = P(x)$$

$$P(y|x) = P(y)$$

## Basic rules

**law of total probability** The *marginal distribution* of a variable is obtained from a joint distribution summing over all possible values of the other variable (*sum rule*)

$$P(x) = \sum_{y \in \mathcal{Y}} P(x, y)$$

$$P(y) = \sum_{x \in \mathcal{X}} P(x, y)$$

**product rule** conditional probability definition implies that

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

## Bayes' rule

$$P(y|x) = \frac{P(x|y)P(y)}{P(x)}$$

## Bayes' rule

### Significance

$$P(y|x) = \frac{P(x|y)P(y)}{P(x)}$$

- allows to “invert” statistical connections between *effect* ( $x$ ) and *cause* ( $y$ ):

$$\text{posterior} = \frac{\text{likelihood} \times \text{prior}}{\text{evidence}}$$

- evidence can be obtained using the sum rule from likelihood and prior:

$$P(x) = \sum_y P(x, y) = \sum_y P(x|y)P(y)$$

## Playing with probabilities

### Use rules!

- Basic rules allow to model a certain probability (e.g. cause given effect) given knowledge of some related ones (e.g. likelihood, prior)
- All our manipulations will be applications of the three basic rules
- Basic rules apply to any number of variables:

$$\begin{aligned}P(y) &= \sum_x \sum_z P(x, y, z) \quad (\text{sum rule}) \\&= \sum_x \sum_z P(y|x, z)P(x, z) \quad (\text{product rule}) \\&= \sum_x \sum_z \frac{P(x|y, z)P(y|z)P(x, z)}{P(x|z)} \quad (\text{Bayes rule})\end{aligned}$$

## Playing with probabilities

### Example

$$\begin{aligned}P(y|x, z) &= \frac{P(x, z|y)P(y)}{P(x, z)} \quad (\text{Bayes rule}) \\&= \frac{P(x, z|y)P(y)}{P(x|z)P(z)} \quad (\text{product rule}) \\&= \frac{P(x|z, y)P(z|y)P(y)}{P(x|z)P(z)} \quad (\text{product rule}) \\&= \frac{P(x|z, y)P(z, y)}{P(x|z)P(z)} \quad (\text{product rule}) \\&= \frac{P(x|z, y)P(y|z)P(z)}{P(x|z)P(z)} \quad (\text{product rule}) \\&= \frac{P(x|z, y)P(y|z)}{P(x|z)}\end{aligned}$$

## Continuous random variables

### Cumulative distribution function

- How to generalize probability mass function to continuous domains?
- Consider probability of *intervals*, e.g.

$$W = (a < X \leq b) \quad A = (X \leq a) \quad B = (X \leq b)$$

- $W$  and  $A$  are mutually exclusive, thus:

$$P(B) = P(A) + P(W) \quad P(W) = P(B) - P(A)$$

- We call  $F(q) = P(X \leq q)$  the *cumulative distribution function* (cdf) of  $X$  (monotonic function)
- The probability of an interval is the difference of two cdf:

$$P(a < X \leq b) = F(b) - F(a)$$

## Continuous random variables

### Probability density function

- The derivative of the cdf is called *probability density function* (pdf):

$$p(x) = \frac{d}{dx}F(x)$$

- The cdf can be computed integrating the pdf:

$$F(q) = P(X \leq q) = \int_{-\infty}^q p(x)dx$$

- Properties:

- $p(x) \geq 0$
- $\int_{-\infty}^{\infty} p(x)dx = 1$

## Continuous random variables

### Note

- The pdf of a value  $x$  can be greater than one, provided the integral is one.
- E.g. let  $p(x)$  be a uniform distribution over  $[a, b]$ :

$$p(x) = \text{Unif}(x; a, b) = \frac{1}{b-a}(a \leq x \leq b)$$

- For  $a = 0$  and  $b = 1/2$ ,  $p(x) = 2$  for all  $x \in [0, 1/2]$  (but the integral is one)

## Properties

### expected value

$$E[x] = \mu = \int_{-\infty}^{\infty} xp(x)dx$$

### variance

$$\text{Var}[x] = \sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 p(x)dx$$

### Note

Definitions and formulas for discrete random variables carry over to continuous random variables with sums replaced by integrals


## Probability distributions

### Gaussian (or normal) distribution

- Bell-shaped curve.
- Parameters:  $\mu$  mean,  $\sigma^2$  variance.
- Probability density function:

$$p(x; \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} \exp -\frac{(x - \mu)^2}{2\sigma^2}$$

- $E[x] = \mu$
- $\text{Var}[x] = \sigma^2$



- Standard normal distribution:  $N(0, 1)$
- Standardization of a normal distribution  $N(\mu, \sigma^2)$

$$z = \frac{x - \mu}{\sigma}$$


## Probability distributions

### Beta distribution

- Defined in the interval  $[0, 1]$
- Parameters:  $\alpha, \beta$
- Probability density function:

$$p(x; \alpha, \beta) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1}(1-x)^{\beta-1}$$





- $E[x] = \frac{\alpha}{\alpha+\beta}$        $\Gamma(x+1) = x\Gamma(x), \Gamma(1) = 1$

- $\text{Var}[x] = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$

*Note*

It models the posterior distribution of parameter  $p$  of a binomial distribution after observing  $\alpha - 1$  independent events with probability  $p$  and  $\beta - 1$  with probability  $1 - p$ .

**Probability distributions**


**Multivariate normal distribution**

- normal distribution for  $d$ -dimensional vectorial data.
- Parameters:  $\mu$  mean vector,  $\Sigma$  covariance matrix.
- Probability density function:

$$p(\mathbf{x}; \mu, \Sigma) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp -\frac{1}{2} (\mathbf{x} - \mu)^T \Sigma^{-1} (\mathbf{x} - \mu)$$

- $E[x] = \mu$
- $\text{Var}[x] = \Sigma$

### Multivariate Normal Distribution



- squared *Mahalanobis distance* from  $\mathbf{x}$  to  $\boldsymbol{\mu}$  is standard measure of distance to mean:


$$r^2 = (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})$$

### Probability distributions

#### Dirichlet distribution

- Defined:  $\mathbf{x} \in [0, 1]^m, \sum_{i=1}^m x_i = 1$
- Parameters:  $\boldsymbol{\alpha} = \alpha_1, \dots, \alpha_m$
- Probability density function:

$$p(x_1, \dots, x_m; \boldsymbol{\alpha}) = \frac{\Gamma(\alpha_0)}{\prod_{i=1}^m \Gamma(\alpha_i)} \prod_{i=1}^m x_i^{\alpha_i - 1}$$



- $E[x_i] = \frac{\alpha_i}{\alpha_0}$       where  $\alpha_0 = \sum_{j=1}^m \alpha_j$
- $\text{Var}[x_i] = \frac{\alpha_i(\alpha_0 - \alpha_i)}{\alpha_0^2(\alpha_0 + 1)}$        $\text{Cov}[x_i, x_j] = \frac{-\alpha_i \alpha_j}{\alpha_0^2(\alpha_0 + 1)}$

*Note*

It models the posterior distribution of parameters  $p$  of a multinomial distribution after observing  $\alpha_i - 1$  times each mutually exclusive event

**Probability laws**

**Expectation of an average**

Consider a sample of  $X_1, \dots, X_n$  i.i.d instances drawn from a distribution with mean  $\mu$  and variance  $\sigma^2$ .

- Consider the random variable  $\bar{X}_n$  measuring the sample average:

$$\bar{X}_n = \frac{X_1 + \dots + X_n}{n}$$

- Its expectation is computed as  $(E[a(X + Y)] = a(E[X] + E[Y]))$ :

$$E[\bar{X}_n] = \frac{1}{n}(E[X_1] + \dots + E[X_n]) = \mu$$

- i.e. the expectation of an average is the true mean of the distribution

### Probability laws

#### variance of an average

- Consider the random variable  $\bar{X}_n$  measuring the sample average:

$$\bar{X}_n = \frac{X_1 + \dots + X_n}{n}$$

- Its variance is computed as  $(\text{Var}[a(X + Y)] = a^2(\text{Var}[X] + \text{Var}[Y])$  for  $X$  and  $Y$  independent):

$$\text{Var}[\bar{X}_n] = \frac{1}{n^2}(\text{Var}[X_1] + \dots + \text{Var}[X_n]) = \frac{\sigma^2}{n}$$

- i.e. the variance of the average *decreases* with the number of observations (the more examples you see, the more likely you are to estimate the correct average)

### Probability laws

#### Chebyshev's inequality

Consider a random variable  $X$  with mean  $\mu$  and variance  $\sigma^2$ .

- Chebyshev's inequality states that for all  $a > 0$ :

$$\Pr[|X - \mu| \geq a] \leq \frac{\sigma^2}{a^2}$$

- Replacing  $a = k\sigma$  for  $k > 0$  we obtain:

$$\Pr[|X - \mu| \geq k\sigma] \leq \frac{1}{k^2}$$

#### Note

Chebyshev's inequality shows that most of the probability mass of a random variable stays within few standard deviations from its mean

### Probability laws

#### The law of large numbers

Consider a sample of  $X_1, \dots, X_n$  i.i.d instances drawn from a distribution with mean  $\mu$  and variance  $\sigma^2$ .

- For any  $\epsilon > 0$ , its sample average  $\bar{X}_n$  obeys:

$$\lim_{n \rightarrow \infty} \Pr[|\bar{X}_n - \mu| > \epsilon] = 0$$

- It can be shown using Chebyshev's inequality and the facts that  $E[\bar{X}_n] = \mu$ ,  $\text{Var}[\bar{X}_n] = \sigma^2/n$ :

$$\Pr[|\bar{X}_n - E[\bar{X}_n]| \geq \epsilon] \leq \frac{\sigma^2}{n\epsilon^2}$$

#### Interpretation

- The accuracy of an empirical statistic increases with the number of samples

## Probability laws

### Central Limit theorem

Consider a sample of  $X_1, \dots, X_n$  i.i.d instances drawn from a distribution with mean  $\mu$  and variance  $\sigma^2$ .

1. Regardless of the distribution of  $X_i$ , for  $n \rightarrow \infty$ , the distribution of the sample average  $\bar{X}_n$  approaches a Normal distribution
2. Its mean approaches  $\mu$  and its variance approaches  $\sigma^2/n$
3. Thus the normalized sample average:

$$z = \frac{\bar{X}_n - \mu}{\frac{\sigma}{\sqrt{n}}}$$

approaches a standard Normal distribution  $N(0, 1)$ .

### Central Limit theorem

#### Interpretation

- The sum of a sufficiently large sample of i.i.d. random measurements is approximately normally distributed
- We don't need to know the form of their distribution (it can be arbitrary)
- Justifies the importance of Normal distribution in real world applications

## Information theory

### Entropy

- Consider a discrete set of symbols  $\mathcal{V} = \{v_1, \dots, v_n\}$  with mutually exclusive probabilities  $P(v_i)$ .
- We aim at designing a binary code for each symbol, minimizing the average length of messages
- Shannon and Weaver (1949) proved that the optimal code assigns to each symbol  $v_i$  a number of bits equal to  
$$-\log P(v_i)$$
- The *entropy* of the set of symbols is the expected length of a message encoding a symbol assuming such optimal coding:

$$H[\mathcal{V}] = E[-\log P(v)] = -\sum_{i=1}^n P(v_i) \log P(v_i)$$

## Information theory

### Cross entropy

- Consider two distributions  $P$  and  $Q$  over variable  $X$
- The *cross entropy* between  $P$  and  $Q$  measures the expected number of bits needed to code a symbol sampled from  $P$  using  $Q$  instead

$$H(P; Q) = E_P[-\log Q(v)] = -\sum_{i=1}^n P(v_i) \log Q(v_i)$$

#### Note

It is often used as a *loss* for binary classification, with  $P$  (empirical) true distribution and  $Q$  (empirical) predicted distribution.

## Information theory

### Relative entropy

- Consider two distributions  $P$  and  $Q$  over variable  $X$
- The *relative entropy* or *Kullback-Leibler (KL) divergence* measures the expected length difference when coding instances sampled from  $P$  using  $Q$  instead:

$$\begin{aligned}D_{KL}(p||q) &= H(P; Q) - H(P) \\ &= - \sum_{i=1}^n P(v_i) \log Q(v_i) + \sum_{i=1}^n P(v_i) \log P(v_i) \\ &= \sum_{i=1}^n P(v_i) \log \frac{P(v_i)}{Q(v_i)}\end{aligned}$$

*Note*

The KL-divergence is not a distance (metric) as it is not necessarily symmetric

## Information theory

### Conditional entropy

- Consider two variables  $V, W$  with (possibly different) distributions  $P$
- The *conditional entropy* is the entropy remaining for variable  $W$  once  $V$  is known:

$$\begin{aligned}H(W|V) &= \sum_v P(v) H(W|V = v) \\ &= - \sum_v P(v) \sum_w P(w|v) \log P(w|v)\end{aligned}$$

## Information theory

### Mutual information

- Consider two variables  $V, W$  with (possibly different) distributions  $P$
- The *mutual information* (or *information gain*) is the reduction in entropy for  $W$  once  $V$  is known:

$$\begin{aligned}I(W; V) &= H(W) - H(W|V) \\ &= - \sum_w p(w) \log p(w) + \sum_v P(v) \sum_w P(w|v) \log P(w|v)\end{aligned}$$

*Note*

It is used e.g. in selecting the best attribute to use in building a decision tree, where  $V$  is the attribute and  $W$  is the label.