

Course “Introduction to SAT & SMT”
TEST

Roberto Sebastiani
DISI, Università di Trento, Italy

June 11th, 2020

769857918

[COPY WITH SOLUTIONS]

1

Let φ be a generic Boolean formula, and let $\varphi_{nnf}^{tree} \stackrel{\text{def}}{=} NNF^{tree}(\varphi)$ and $\varphi_{nnf}^{dag} \stackrel{\text{def}}{=} NNF^{dag}(\varphi)$, s.c. $NNF()^{tree}$ and $NNF()^{dag}$ are the conversion into negative normal form using a tree and a DAG representation of the formulas respectively.

Let $|\varphi|$, $|\varphi_{nnf}^{tree}|$ and $|\varphi_{nnf}^{dag}|$ denote the size of φ , φ_{nnf}^{tree} and φ_{nnf}^{dag} respectively.

For each of the following sentences, say if it is true or false.

- (a) $|\varphi_{nnf}^{tree}|$ is in worst-case polynomial in size wrt. $|\varphi|$. [Solution: False.]
- (b) $|\varphi_{nnf}^{dag}|$ is in worst-case polynomial in size wrt. $|\varphi|$. [Solution: True.]
- (c) φ_{nnf}^{dag} has the same number of distinct Boolean variables as φ has. [Solution: True.]
- (d) A model for φ_{nnf}^{dag} (if any) is also a model for φ , and vice versa. [Solution: True.]


2

Using the variable ordering “ A_1, A_2, A_3, A_4 ”, draw the OBDD corresponding to the following formulas:

$$A_1 \wedge (\neg A_1 \vee \neg A_2) \wedge (A_2 \vee A_3) \wedge (\neg A_3 \vee A_4)$$

[Solution:

SOLUTION: The formula is represented by the following OBDD:



]

4

Consider the following piece of a much bigger formula, which has been fed to a CDCL SAT solver:

$$\begin{aligned}
 c_1 &: \neg A_7 \vee A_2 \\
 c_2 &: A_4 \vee A_1 \vee A_{11} \\
 c_3 &: A_8 \vee \neg A_6 \vee \neg A_4 \\
 c_4 &: \neg A_5 \vee \neg A_1 \\
 c_5 &: A_7 \vee \neg A_8 \\
 c_6 &: A_7 \vee A_6 \vee A_9 \\
 c_7 &: \neg A_7 \vee A_3 \vee \neg A_{12} \\
 c_8 &: A_4 \vee A_5 \vee A_{10} \\
 &\dots
 \end{aligned}$$

Suppose the solver has decided, in order, the following literals (possibly interleaved by others not occurring in the above clauses):

$$\{\dots, \neg A_9, \dots, \neg A_{10}, \dots, \neg A_{11}, \dots, A_{12}, \dots, A_{13}, \dots, \neg A_7\}$$

- (a) List the sequence of unit-propagations following after the last decision, each literal tagged (in square brackets) by its antecedent clause

[Solution:

$$\begin{array}{l}
 \neg A_8 \quad [c_5] \\
 A_6 \quad [c_6] \\
 \neg A_4 \quad [c_3] \\
 A_5 \quad [c_8] \\
 A_1 \quad [c_2]
 \end{array}$$

]

- (b) Derive the conflict clause via conflict analysis by means of the 1st-UIP technique

[Solution:

$$\frac{
 \frac{
 A_4 \vee A_5 \vee A_{10} \quad \frac{
 A_4 \vee A_1 \vee A_{11} \quad \overbrace{\neg A_5 \vee \neg A_1}^{\text{Conflicting cl.}}
 }{
 A_4 \vee \neg A_5 \vee A_{11} \quad (A_1)
 }
 }{
 A_4 \vee A_5 \vee A_{10} \quad (A_5)
 }
 }{
 \underbrace{A_4 \vee A_{10} \vee A_{11}}_{\text{1st UIP}}
 }$$

]

- (c) Using the 1st-UIP backjumping strategy, update the list of literals above after the backjumping step and the unit-propagation of the UIP

[Solution: $\{\dots, \neg A_9, \dots, \neg A_{10}, \dots, \neg A_{11}, A_4\}$]

5

Consider the following CNF formula:

$$\begin{aligned}
 & (A_7) \wedge \\
 & (A_8 \vee \neg A_7) \wedge \\
 & (\neg A_4 \vee \neg A_7 \vee \neg A_5) \wedge \\
 & (\neg A_8 \vee \neg A_6 \vee A_1) \wedge \\
 & (A_2 \vee \neg A_7 \vee \neg A_6) \wedge \\
 & (\neg A_8 \vee A_6 \vee \neg A_2) \wedge \\
 & (\neg A_1 \vee \neg A_5 \vee \neg A_2) \wedge \\
 & (A_2 \vee \neg A_6 \vee \neg A_8) \wedge \\
 & (\neg A_1 \vee \neg A_5 \vee A_8) \wedge \\
 & (A_2 \vee A_7 \vee \neg A_6) \wedge \\
 & (\neg A_6 \vee A_4 \vee \neg A_6) \wedge \\
 & (A_3 \vee A_8 \vee \neg A_7)
 \end{aligned}$$

Decide *quickly* if it is satisfiable or not, and briefly explain why.

[Solution: After unit-propagating A_7, A_8 :

$$\begin{aligned}
 & (A_7) \wedge \\
 & (A_8) \wedge \\
 & (\neg A_4 \vee \neg A_5) \wedge \\
 & (\neg A_6 \vee A_1) \wedge \\
 & (A_2 \vee \neg A_6) \wedge \\
 & (A_6 \vee \neg A_2) \wedge \\
 & (\neg A_1 \vee \neg A_5 \vee \neg A_2) \wedge \\
 & (A_2 \vee \neg A_6) \wedge \\
 & (\neg A_1 \vee \neg A_5 \vee A_8) \wedge \\
 & (A_2 \vee A_7 \vee \neg A_6) \wedge \\
 & (\neg A_6 \vee A_4 \vee \neg A_6) \wedge \\
 & (A_3 \vee A_8)
 \end{aligned}$$

the result is a Horn formula with no positive unit clauses.

Therefore, the assignment $\{\neg A_i\}_{i=1}^8$ is a model.]

6

Consider the following Boolean formulas:

$$\varphi_1 \stackrel{\text{def}}{=} \begin{array}{l} (\neg A_7 \vee \neg A_3) \wedge \\ (A_7 \vee \neg A_3) \wedge \\ (A_2) \wedge \\ (\neg A_2 \vee \neg A_4) \end{array}$$

$$\varphi_2 \stackrel{\text{def}}{=} \begin{array}{l} (A_3 \vee A_5) \wedge \\ (A_4 \vee \neg A_1) \wedge \\ (\neg A_5 \vee A_1) \end{array}$$

which are such that $\varphi_1 \wedge \varphi_2 \models \perp$. For each of the following formulas, say if it is a Craig interpolant for (φ_1, φ_2) or not.

[Solution: Recall that a Craig interpolant for (φ_1, φ_2) s.t. $\varphi_1 \wedge \varphi_2 \models \perp$ is a formula ψ s.t.

1. $\varphi_1 \models \psi$
2. $\psi \wedge \varphi_2 \models \perp$
3. all atoms in ψ occur in both φ_1 and φ_2 .

]

(a)

$$\begin{array}{l} (\neg A_7 \vee \neg A_3) \wedge \\ (A_7 \vee \neg A_3) \wedge \\ (\neg A_4) \end{array}$$

[Solution: No, it is not a solution because it does not verify condition 3.]

(b)

$$(\neg A_4)$$

[Solution: No, it is not a solution because it does not verify condition 2.]

(c)

$$(\neg A_3 \wedge \neg A_4)$$

[Solution: Yes]

7

Consider the following formula in the theory \mathcal{EUF} of linear arithmetic on the Rationals.

$$\begin{aligned} \varphi = & \{(f(x) = f(f(y))) \vee A_2\} \wedge \\ & \{\neg(h(x, f(y)) = h(g(x), y)) \vee \underline{\neg(h(x, g(z) = h(f(x), y)))} \vee \neg A_1\} \wedge \\ & \{A_1 \vee (h(x, y) = h(y, x))\} \wedge \\ & \{(x = \underline{f(x)}) \vee A_3 \vee \neg A_1\} \wedge \\ & \{\underline{\neg(w(x) = g(f(y)))} \vee A_1\} \wedge \\ & \{\underline{\neg A_2} \vee (w(g(x)) = w(f(x)))\} \wedge \\ & \{A_1 \vee \underline{(y = g(z))} \vee A_2\} \end{aligned}$$

and consider the partial truth assignment μ given by the underlined literals above:

$$\{\neg(w(x) = g(f(y))), \neg A_2, \neg(h(x, g(z) = h(f(x), y))), (x = f(x)), (y = g(z))\}.$$

1. Does (the Boolean abstraction of) μ propositionally satisfy (the Boolean abstraction of) φ ? [**Solution:** No, since there are two clauses which are not satisfied.]

2. Is μ satisfiable in \mathcal{EUF} ?

- (a) If no, find a minimal conflict set for μ and the corresponding conflict clause C .
- (b) If yes, show one unassigned literal which can be deduced from μ , and show the corresponding deduction clause C .

[**Solution:**

No, because it contains the following \mathcal{EUF} conflict set:

$$\{\neg(h(x, g(z) = h(f(x), y))), (x = f(x)), (y = g(z))\},$$

which corresponds to the following conflict clause:

$$(h(x, g(z) = h(f(x), y))) \vee \neg(x = f(x)), \neg(y = g(z))$$

]

8

Consider the following set of clauses φ in the theory of linear arithmetic on the Integers \mathcal{EUF} .

$$\left\{ \begin{array}{l} (\neg(x = y) \vee (f(x) = f(y))), \\ (\neg(x = y) \vee \neg(f(x) = f(y))), \\ ((x = y) \vee (f(x) = f(y))), \\ ((x = y) \vee \neg(f(x) = f(y))) \end{array} \right\}$$

Say which of the following sets is a \mathcal{EUF} -unsatisfiable core of φ and which is not. For each one, explain why.

(a)

$$\left\{ \begin{array}{l} (\neg(x = y) \vee \neg(f(x) = f(y))), \\ ((x = y) \vee (f(x) = f(y))), \\ ((x = y) \vee \neg(f(x) = f(y))) \end{array} \right\}$$

[Solution: yes, because it is a subset of φ and it is inconsistent in \mathcal{EUF} .]

(b)

$$\left\{ \begin{array}{l} (\neg(x = y) \vee (f(x) = f(y))), \\ ((x = y) \vee (f(x) = f(y))), \\ ((x = y) \vee \neg(f(x) = f(y))) \end{array} \right\}$$

[Solution: no, because is not inconsistent in \mathcal{EUF} (e.g., $\{ (x = y), (f(x) = f(y)) \}$ is \mathcal{EUF} -consistent solution).]

(c)

$$\left\{ \begin{array}{l} (\neg(x = y) \vee \neg(f(x) = f(y))), \\ ((x = y) \vee (f(x) = f(y))), \\ ((x = y) \vee \neg(f(x) = f(y))), \\ ((x = f(y))) \end{array} \right\}$$

[Solution: no, because it is not a subset of φ .]

9

Let \mathcal{LRA} be the logic of linear arithmetic over the rationals and \mathcal{EUF} be the logic of equality and uninterpreted functions. Consider the following pure formula φ in the combined logic $\mathcal{LRA} \cup \mathcal{EUF}$:

$$(x = 1.0) \wedge (h = 1.0) \wedge (k = 1.0) \wedge (y = 2h - k) \wedge (z < w) \quad (1)$$

$$(z = f(x)) \wedge (w = f(y)) \quad (2)$$

Say which variables are interface variables, list the interface equalities for this formula (modulo symmetry), and decide whether this formulas is $\mathcal{LRA} \cup \mathcal{EUF}$ -satisfiable or not, using either Nelson-Oppen or Delayed Theory Combination.

[Solution: Only x, y, z, w occur both in \mathcal{LRA} -atoms (2) and in \mathcal{EUF} -atoms (1). Thus x, y, z, w are the interface variables, and $x = y, x = z, x = w, y = z, y = w, z = w$ are the interface equalities.

Nelson-Oppen: From (2) in \mathcal{LRA} we infer the interface equality $(x = y)$. Adding the latter to (1), we infer the interface equality $(z = w)$ in \mathcal{EUF} . Adding the latter to (2), we get a contradiction in \mathcal{LRA} against $(z < w)$.

Delayed Theory Combination: By unit-propagation, φ causes only one branch containing all its literals. Then the SAT solver assigns first a negative value to the interface equality $(x=y)$, adding $\neg(x = y)$ to the assignment, which is found inconsistent in \mathcal{LRA} :

$$(h = 1.0) \wedge (k = 1.0) \wedge (x = 1.0) \wedge (y = 2h - k) \wedge \neg(x = y). \quad (3)$$

Then the SAT solver backtracks, adding $(x = y)$ to the assignment. Then the SAT solver assigns first a negative value to the interface equality $(z=w)$, adding $\neg(z = w)$ to the assignment, which is found inconsistent in \mathcal{LRA} :

$$(z = f(x)) \wedge (w = f(y)) \wedge (x = y) \wedge \neg(z = w). \quad (4)$$

Thus, with either technique, we can conclude that φ is $\mathcal{LRA} \cup \mathcal{EUF}$ -unsatisfiable.]

10

Consider the following formulas in difference logic (\mathcal{DL}):

$$\varphi_1 \stackrel{\text{def}}{=} \begin{aligned} &(x_2 - x_3 \leq -4) \wedge \\ &(x_3 - x_4 \leq -6) \wedge \\ &(x_5 - x_6 \leq 4) \wedge \\ &(x_6 - x_1 \leq 2) \wedge \\ &(x_6 - x_7 \leq -2) \wedge \\ &(x_7 - x_8 \leq 1) \end{aligned}$$

$$\varphi_2 \stackrel{\text{def}}{=} \begin{aligned} &(x_4 - x_9 \leq 2) \wedge \\ &(x_9 - x_5 \leq 0) \wedge \\ &(x_1 - x_2 \leq 1) \end{aligned}$$

which are such that $\varphi_1 \wedge \varphi_2 \models_{\mathcal{DL}} \perp$. For each of the following formulas, say if it is a Craig interpolant in \mathcal{DL} for (φ_1, φ_2) , and explain why.

[Solution: Recall that a Craig interpolant for (φ_1, φ_2) s.t. $\varphi_1 \wedge \varphi_2 \models_{\mathcal{DL}} \perp$ is a formula ψ s.t.

1. $\varphi_1 \models_{\mathcal{DL}} \psi$
2. $\psi \wedge \varphi_2 \models_{\mathcal{DL}} \perp$
3. all symbols in ψ occur in both φ_1 and φ_2 .

]

(a) $(x_2 - x_3 + x_6 - x_1 \leq -2)$

[Solution: no, because, e.g., x_3 is not a symbol occurring in φ_2 .]

(b) $(x_2 - x_4 \leq -10)$

[Solution: No, because it violates condition 2.]

(c) $(x_2 - x_4 \leq -10) \wedge$
 $(x_5 - x_1 \leq 6)$

[Solution: yes, because it is a \mathcal{DL} formula and it verifies all conditions 1., 2., 3.]