Computability Assignment Year 2012/13 - Number 6

Please keep this file anonymous: do not write your name inside this file.

 $More \ information \ about \ assignments \ at \ http://disi.unitn.it/\sim zunino/teaching/computability/assignments$

Please do not submit a file containing only the answers; edit this file, instead, filling the answer sections.

1 Question

Write a λ -term M implementing the following specification:

$$M \, \lceil n \rceil = \lceil \lambda x_0 \dots x_n \dots x_n x_{n-1} \dots x_0 \rceil$$

(Note: The notation $\lceil n \rceil$ above stands for the numeral n, while $\lceil N \rceil$ stands for $\lceil \# N \rceil$ – inside L_YX it's hard to tell them apart, but will appear correctly in the PDFs)

1.1 Answer

Write your answer here.

2 Question

Write a λ -term M which, when given as input $\lceil N \rceil$, evaluates to $\lceil O \rceil$, where O is obtained from N by replacing every syntactic occurrence of Ω with \mathbf{I} .

To the purpose of this exercise, assume $\Omega = (\lambda x_0.x_0x_0)(\lambda x_0.x_0x_0)$ and $\mathbf{I} = \lambda x_0.x_0$.

For example, here are some expected outputs:

$$\begin{split} M & \lceil \lambda x_5.\Omega \rceil =_{\beta\eta} \lceil \lambda x_5.\mathbf{I} \rceil \\ M & \lceil \lambda x_3.\mathbf{K}\Omega \rceil =_{\beta\eta} \lceil \lambda x_3.\mathbf{K}\mathbf{I} \rceil \\ M & \lceil \lambda x_1.x_1\Omega(\lambda x_7.x_1\Omega) \rceil =_{\beta\eta} \lceil \lambda x_1.x_1\mathbf{I}(\lambda x_7.x_1\mathbf{I}) \rceil \\ M & \lceil (\lambda x_0.x_0x_0)(\lambda x_0.x_0x_0) \rceil =_{\beta\eta} \lceil \mathbf{I} \rceil \\ M & \lceil (\lambda x_1.x_1x_1)(\lambda x_1.x_1x_1) \rceil =_{\beta\eta} \lceil (\lambda x_1.x_1x_1)(\lambda x_1.x_1x_1) \rceil \end{split}$$

Hint: use **Sd**, etc. as appropriate.

2.1 Answer

Write your answer here.