

# Computability Assignment

## Year 2012/13 - Number 1

Please keep this file anonymous: do not write your name inside this file.

More information about assignments at <http://disi.unitn.it/~zunino/teaching/computability/assignments>

### 1 Question

Define a binary property  $p(x, y)$  over natural numbers such that we have both

1.  $\forall x \in \mathbb{N}. \exists y \in \mathbb{N}. p(x, y) \iff \neg \exists x \in \mathbb{N}. \forall y \in \mathbb{N}. \neg p(x, y)$
2.  $\neg \exists y \in \mathbb{N}. \forall x \in \mathbb{N}. p(x, y) \iff \forall y \in \mathbb{N}. \exists x \in \mathbb{N}. \neg p(x, y)$

Provide a definition for  $p$ , and a proof for the above claims.

#### 1.1 First Trial:

Let  $\langle x, y \rangle \in p \iff y = f(x)$ , where  $f$  is a recursive function so defined:

$$f(x) = \begin{cases} y = x/2 & \text{if } \exists! n \in \mathbb{N}. x = 2 * n \\ y = (x - 1)/2 & \text{if } \exists! n \in \mathbb{N}. x = 2 * n + 1 \end{cases}, \text{ where it is implicit that}$$

$y \in \mathbb{N}$  and that the symbols  $+$ ,  $-$ ,  $/$  follow the usual arithmetic semantics among natural numbers.

STATEMENT 1: (PROOF BY ENUMERATION OF CASES)

Let  $x_0 \in \mathbb{N}$  be an arbitrarily chosen number, there are two possible cases:  $x_0$  may be either even or odd.

CASE A:  $x_0$  is even.

If  $x_0$  is even, there exists by definition a (unique)  $n \in \mathbb{N}$  such that  $x_0 = 2 * n$ .

Therefore by the actual definition of function  $f$ , we have that  $y_0 = f(x_0) = x_0/2$ , and being more precise  $y_0 = f(x_0) = f(2 * n) = 2 * n/2 = n$ .

Since  $y_0 = n \in \mathbb{N}$ , the tuple  $\langle x_0, y_0 \rangle$  belongs to the binary relationship  $p$ .

CASE B:  $x_0$  is odd.

If  $x_0$  is odd, there exists by definition a (unique)  $n \in \mathbb{N}$  such that  $x_0 = 2 * n + 1$ .

Therefore by the actual definition of function  $f$ , we have that  $y_0 = f(x_0) = (x_0 - 1)/2$  which means  $y_0 = f(x_0) = f(2 * n + 1) = ((2 * n + 1) - 1)/2 = (2 * n)/2 = n$ .

Since  $y_0 = n \in \mathbb{N}$ , the tuple  $\langle x_0, y_0 \rangle$  belongs to the binary relationship  $p$ .

CONCLUSION:

Having chosen  $x_0 \in \mathbb{N}$  arbitrarily and examined all the possible cases, we may generalize and say that for each  $x \in \mathbb{N}$  there exists  $y \in \mathbb{N}$  such that  $p(x, y)$ , which is precisely what Statement 1 is standing for.

STATEMENT 2:

Let  $y \in \mathbb{N}$  be, for absurdum, a chosen number such that  $\forall x \in \mathbb{N}. p(x, y)$ .

Since the property holds for all  $x \in \mathbb{N}$ , it certainly has to hold for a strict subset of  $\mathbb{N}$ , let's say  $A_m = \{x | x = m \vee x = s(m) \vee x = s(s(m))\}$  ( $s(x)$  is the usual successor function).

We may take an arbitrary set  $A_m = \{m, m + 1, m + 2\}$ , hence we make no restriction and use it directly without instantiating  $m$ .

For hypothesis  $\langle m, y \rangle \in p, \langle m + 1, y \rangle \in p$  and  $\langle m + 2, y \rangle \in p$ .

Our goal is to show that the following statement  $y = f(m) = f(m + 1) = f(m + 2)$  is false, thus falsifying the entire conjecture.

There are two cases:

$$\text{if } m = 2 * k \text{ then } \begin{cases} f(m) = f(2 * k) = 2 * k / 2 & = k \\ f(m + 1) = f(2 * k + 1) = (2 * k + 1 - 1) / 2 & = k \\ f(m + 2) = f(2 * k + 2) = (2 * k + 2) / 2 & = k + 1 \end{cases},$$

which clearly contradicts the hypothesis.

$$\text{if } m = 2 * k + 1 \text{ then } \begin{cases} f(m) = f(2 * k + 1) = (2 * k + 1 - 1) / 2 & = k \\ f(m + 1) = f(2 * k + 2) = (2 * k + 2) / 2 & = k + 1, \\ f(m + 2) = f(2 * k + 3) = (2 * k + 3 - 1) / 2 & = k + 1 \end{cases},$$

which clearly contradicts the hypothesis.

Therefore there exists no  $y \in \mathbb{N}$  with such properties, therefore we can discard the initial hypothesis and conclude  $\neg \exists y \in \mathbb{N}. \forall x \in \mathbb{N}. p(x, y)$ .

## 1.2 Second Trial.

Let  $\langle x, y \rangle \in p \iff y = s(x)$ , so that  $p$  may be read as the property relationship of “ $x$  is the predecessor of  $y$ ” or, conversely, “ $y$  is the successor of  $x$ ”. The function  $s(x)$  is total and injective, while the property  $p$  is anti-reflexive, anti-symmetric and clearly not transitive [proof by exercise].

1:

The first statement is straightforward, since  $\mathbb{N}$  is a well-founded set that may be recursively defined as

$$0 \in \mathbb{N} \text{ and } \forall x \in \mathbb{N} \Rightarrow s(x) \in \mathbb{N}$$

then it is obvious that  $\forall x \in \mathbb{N}$ , its successor  $s(x)$  belongs to the set of  $\mathbb{N}$ .

2:

The second statement is obviously right, since otherwise it would violate both the injective property of  $s(x)$  and the anti-reflexive and anti-symmetric properties of  $p$ .

Let's try to sort out a proof:

Let  $y \in \mathbb{N}$  be, for absurdum, a chosen number such that  $\forall x \in \mathbb{N}. p(x, y)$ .

Since  $y \in \mathbb{N}$ , by the property of  $\mathbb{N}$  there exist a (possibly large) finite index  $n \in \mathbb{N}$  such that  $y = s_n(s_{n-1}(\dots s_1(0)\dots))$ . Also, since  $y \in \mathbb{N}$ , there must exist  $y' \in \mathbb{N}$  such that  $y' = s(y)$ , which means  $\langle y, y' \rangle \in p$ .

For hypothesis  $y$  is such that  $\forall x \in \mathbb{N}. p(x, y)$ , therefore also  $\langle y', y \rangle \in p$  and  $y = s(y')$ .

Hence one could deduct  $y = s(y') = s(s(y))$ , which follows into:

$$y = s_n(s_{n-1}(\dots s_1(0)\dots)) = s_{n+2}(s_{n+1}(\dots s_1(0)\dots))) = s(s(y))$$

This is clearly possible only if  $n = n + 2$ , but clearly  $0 \neq 2$ .

Since the only hypothesis made was the existence of a  $y \in \mathbb{N}$  with such satisfying the second statement, and it was shown that the property didn't hold for an arbitrarily chosen  $y$ , we can discard the claim and deduce  $\neg \exists y \in \mathbb{N}. \forall x \in \mathbb{N}. p(x, y)$  is true.

Viger