# Computability Assignment Year 2012/13 - Number 2

Please keep this file anonymous: do not write your name inside this file. More information about assignments at http://disi.unitn.it/~zunino/teaching/computability/assignments

## 1 Question

Let A, B be two sets. Prove that the properties below are equivalent.

- $A = \emptyset \lor B = \emptyset$
- $\bullet \ A \times B = \emptyset$

#### 1.1 Answer

The Result of the cartesian product between 2 empty sets is empty because cannot be formed order pair,

for e.g:

- $A \times B = \{ \emptyset \{ \emptyset \} \} = \emptyset$ ; according with the definition of cartesian product which states  $A \times B = \{ \langle a, b \rangle | a \epsilon A \land b \epsilon B \}$
- $a \notin A \land b \notin B = \emptyset \iff \neg (a \in A) \land \neg (b \in B) = \emptyset$  $\neg \exists a(a \in A) \land \neg \exists b(b \in B) = \emptyset$

### 2 Preliminaries

Given an infinite sequence of sets  $(A_i)_{i\in\mathbb{N}}$ , we define  $\bigcup_{i=0}^{\infty} A_i = \bigcup \{A_i \mid i \in \mathbb{N}\}$ and  $\bigcup_{i=0}^k A_i = \bigcup \{A_i \mid i \in \mathbb{N} \land i \leq k\} = A_0 \cup A_1 \cup \cdots \cup A_k$ .

## 3 Question

Assume  $(A_i)_{i\in\mathbb{N}}$  to be an infinite sequence of sets of natural numbers, satisfying

$$A_0 \subseteq A_1 \subseteq A_2 \subseteq A_3 \cdots \subseteq \mathbb{N} \ (*)$$

For each property  $p_i$  shown below, state whether

- the hypothesis (\*) is sufficient to conclude that  $p_i$  holds; or
- the hypothesis (\*) is sufficient to conclude that  $p_i$  does not hold; or
- the hypothesis (\*) is not sufficient to conclude anything about the truth of  $p_i$ .

Justify your answers (briefly).

- 1.  $p_1: \forall k \in \mathbb{N}. A_k = \bigcup_{i=0}^k A_i$
- 2.  $p_2$ : for all *i*, if  $A_i$  is infinite, then  $A_i = A_{i+1}$
- 3.  $p_3$ : if  $\forall i \in \mathbb{N}$ .  $A_i \neq A_{i+1}$ , then  $\bigcup_{i=0}^{\infty} A_i = \mathbb{N}$
- 4.  $p_4$ : if  $\forall i \in \mathbb{N}$ .  $A_i$  is finite, then  $\bigcup_{i=0}^{\infty} A_i$  is finite
- 5.  $p_5$ : if  $\forall i \in \mathbb{N}$ .  $A_i$  is finite, then  $\bigcup_{i=0}^{\infty} A_i$  is infinite
- 6.  $p_6$ : if  $\forall i \in \mathbb{N}$ .  $A_i$  is infinite, then  $\bigcup_{i=0}^{\infty} A_i$  is infinite

#### 3.1 Answer

1.  $p_1: \forall k \in \mathbb{N}.A_k = \bigcup_{i=0}^k A_i$ - the hypothesis (\*) is sufficient to conclude that  $p_i$  holds;

- $A_k = \{0, 1, 2, ..., k\} / /$  this is a Set of sets.
- $A_i = \{x : \forall i \in A_k \ (x \le k)\}$

