

# Computability Assignment

## Year 2012/13 - Number 4

Please keep this file anonymous: do not write your name inside this file.

More information about assignments at <http://disi.unitn.it/~zunino/teaching/computability/assignments>

**Please do not submit a file containing only the answers; edit this file, instead, filling the answer sections.**

## 1 Preliminaries

A partial function  $g$  is said to be a *restriction* of a partial function  $f$ , written  $g \subseteq f$  iff

$$\forall x \in \text{dom}(g). g(x) = f(x)$$

Note: this notation “overloads” the symbol  $\subseteq$ . Indeed, we shall write  $A \subseteq B$  to express a subset relation between two sets, and  $g \subseteq f$  to express a restriction relation between two functions.

(From a formal point of view, since we defined functions as set of pairs the two notions coincide: the restriction relation above is equivalent to requiring that  $\langle a, b \rangle \in g \implies \langle a, b \rangle \in f$  for all  $a, b$ , which indeed states that  $g$  is a “subset” of  $f$ ).

## 2 Question

Let  $\mathcal{F}$  be the set of partial functions  $\{f \in (\mathbb{N} \rightsquigarrow \mathbb{N}) \mid \forall x \in \mathbb{N}. f(2 \cdot x) = x\}$ .

- Define two distinct partial functions  $f_1, f_2$  which belong to  $\mathcal{F}$ . (I.e, provide two such examples.)
- Define two distinct partial functions  $g_1, g_2$  which do *not* belong to  $\mathcal{F}$ . (I.e, provide two such examples.)
- Define a partial function  $f \in \mathcal{F}$ , and consider the set of its *finite* restrictions  $\mathcal{G} = \{g \in (\mathbb{N} \rightsquigarrow \mathbb{N}) \mid g \subseteq f \wedge \text{dom}(g) \text{ finite}\}$ .
  - Define two distinct partial functions  $h_1, h_2$  which belong to  $\mathcal{G}$ . (I.e, provide two such examples.)

- Prove whether  $\mathcal{F} \cap \mathcal{G} = \emptyset$ .

## 2.1 Answer

- $f_1(x) = \begin{cases} \frac{x}{2} & \text{if } x \text{ is even} \\ \text{undefined} & \text{o.w.} \end{cases}$

- $f_2(4.x) = \begin{cases} 2x & \text{if } x \geq 10 \\ \text{undefined} & \text{o.w.} \end{cases}$

(RZ: avoid this kind of definition, what should  $f_2(5)$  be, for instance? It's not clear to me.)

- $g_1(x+1) = \begin{cases} x+1 & \text{if } x \text{ is odd} \\ \text{undefined} & \text{o.w.} \end{cases}$

- $g_2(x) = \begin{cases} 2x+1 & \text{if } x \text{ is even} \\ \text{undefined} & \text{o.w.} \end{cases}$

- I consider the above function  $f = f_1 \in \mathcal{F}$ .

- $h_1(x) = \begin{cases} \frac{x}{2} & x \leq 100 \\ \text{undefined} & \text{o.w.} \end{cases}$

- $h_2(6.x) = \begin{cases} 3x & x \leq 200 \\ \text{undefined} & \text{o.w.} \end{cases}$

- $\mathcal{F} \cap \mathcal{G} = \emptyset$ . since in  $\mathcal{F}$ , the domain of each function is infinite, as it has to include all the even numbers. While in  $\mathcal{G}$ , all functions are defined on a finite domain.

## Note.

The next part is an advanced exercise. I'd suggest to **skip** it, unless you want an extra challenge.

## 3 Preliminaries

Let  $\mathcal{R}$  be a set of inference rules over elements of a set  $A$ . Then,  $\mathcal{R}$  induces a function  $\hat{\mathcal{R}} \in (\mathcal{P}(A) \rightarrow \mathcal{P}(A))$  given by

$$\hat{\mathcal{R}}(X) = \{y \mid \exists (\frac{x_1 \cdots x_n}{z}) \in \mathcal{R} \wedge y = z \wedge \forall i \in \{1, \dots, n\}. x_i \in X\}$$

## 4 Question

Let  $m, n$  range over natural numbers. Consider the following set of inference rules  $\mathcal{R}$

$$\frac{n \ m}{n \cdot m} \quad \frac{}{1} \quad \frac{n}{n \cdot 2}$$

and the sets

$$E = \{2 \cdot n \mid n \in \mathbb{N}\} \quad O = \{2 \cdot n + 1 \mid n \in \mathbb{N}\}$$

Then, answer the questions below.

1. State whether  $\hat{\mathcal{R}}(O) \subseteq O$
2. State whether  $O \subseteq \hat{\mathcal{R}}(O)$
3. State whether  $\hat{\mathcal{R}}(E) \subseteq E$
4. State whether  $E \subseteq \hat{\mathcal{R}}(E)$
5. State whether  $\hat{\mathcal{R}}(\mathbb{N}) \subseteq \mathbb{N}$
6. State whether  $\mathbb{N} \subseteq \hat{\mathcal{R}}(\mathbb{N})$
7. State whether  $\hat{\mathcal{R}}(E \cup \{1\}) \subseteq E \cup \{1\}$

You may wish to exploit the answer for some question when answering another. Finally:

1. Characterize the minimum fixed point of  $\hat{\mathcal{R}}$ , i.e.  $\bigcap \{X \mid \hat{\mathcal{R}}(X) = X\}$
2. Characterize the maximum fixed point of  $\hat{\mathcal{R}}$ , i.e.  $\bigcup \{X \mid \hat{\mathcal{R}}(X) = X\}$

### 4.1 Answer