## Computability Assignment Year 2012/13 - Number 4

Please keep this file anonymous: do not write your name inside this file.

More information about assignments at http://disi.unitn.it/~zunino/teaching/computability/assignments Please do not submit a file containing only the answers; edit this file, instead, filling the answer sections.

### 1 Preliminaries

A partial function g is said to be a restriction of a partial function f , written  $g\subseteq f$  iff

$$\forall x \in \mathsf{dom}(g). \ g(x) = f(x)$$

Note: this notation "overloads" the symbol  $\subseteq$ . Indeed, we shall write  $A \subseteq B$  to express a subset relation between two sets, and  $g \subseteq f$  to express a restriction relation between two functions.

(From a formal point of view, since we defined functions as set of pairs the two notions coincide: the restriction relation above is equivalent to requiring that  $\langle a, b \rangle \in g \implies \langle a, b \rangle \in f$  for all a, b, which indeed states that g is a "subset" of f).

## 2 Question

Let  $\mathcal{F}$  be the set of partial functions  $\{f \in (\mathbb{N} \rightsquigarrow \mathbb{N}) | \forall x \in \mathbb{N}. f(2 \cdot x) = x\}$ .

- Define two distinct partial functions  $f_1, f_2$  which belong to  $\mathcal{F}$ . (I.e, provide two such examples.)
- Define two distinct partial functions  $g_1, g_2$  which do *not* belong to  $\mathcal{F}$ . (I.e, provide two such examples.)
- Define a partial function f ∈ F, and consider the set of its finite restrictions G = {g ∈ (N → N)|g ⊆ f ∧ dom(g) finite}.
  - Define two distinct partial functions  $h_1, h_2$  which belong to  $\mathcal{G}$ . (I.e, provide two such examples.)

- Prove whether  $\mathcal{F} \cap \mathcal{G} = \emptyset$ .

#### 2.1 Answer

• 
$$f_1(z) = \begin{cases} 0, & z = 0; \\ undefined, & otherwise. \end{cases}$$
;  $f_2(z) = \begin{cases} 0, & z = 0; \\ 1, & z = 2; \\ undefined, & otherwise. \end{cases}$ 

.

(RZ: they do not belong to  $\mathcal{F}$ )

• 
$$g_1(z) = \begin{cases} 1, & z = 0; \\ undefined, & otherwise. \end{cases}; g_2(z) = \begin{cases} 1, & z = 0; \\ 2, & z = 2; \\ undefined, & otherwise. \end{cases}$$
  
(RZ: ok)

•  $\mathcal{F} \cap \mathcal{G} = \emptyset \iff \neg \exists f. f \in \mathcal{F} \land f \in \mathcal{G}$ , but  $h_1 \in \mathcal{G} \land h_1 \in \mathcal{F} \Longrightarrow \mathcal{F} \cap \mathcal{G} \neq \emptyset$ ??? (RZ: What is  $h_1$ ?)

# Note.

The next part is an advanced exercise. I'd suggest to  ${\bf skip}$  it, unless you want an extra challenge.

### 3 Preliminaries

Let  $\mathcal{R}$  be a set of inference rules over elements of a set A. Then,  $\mathcal{R}$  induces a function  $\hat{\mathcal{R}} \in (\mathcal{P}(A) \to \mathcal{P}(A))$  given by

$$\hat{\mathcal{R}}(X) = \{ y \mid \exists (\frac{x_1 \dots x_n}{z}) \in \mathcal{R} \land y = z \land \forall i \in \{1, \dots, n\} . x_i \in X \}$$

### 4 Question

Let m,n range over natural numbers. Consider the following set of inference rules  $\mathcal R$ 

$$\frac{n \ m}{n \cdot m} \qquad \frac{1}{1} \qquad \frac{n}{n \cdot 2}$$

an the sets

$$E = \{2 \cdot n \mid n \in \mathbb{N}\} \qquad O = \{2 \cdot n + 1 \mid n \in \mathbb{N}\}$$

Then, answer the questions below.

- 1. State whether  $\hat{\mathcal{R}}(O) \subseteq O$
- 2. State whether  $O \subseteq \hat{\mathcal{R}}(O)$

- 3. State whether  $\hat{\mathcal{R}}(E) \subseteq E$
- 4. State whether  $E \subseteq \hat{\mathcal{R}}(E)$
- 5. State whether  $\hat{\mathcal{R}}(\mathbb{N}) \subseteq \mathbb{N}$
- 6. State whether  $\mathbb{N} \subseteq \hat{\mathcal{R}}(\mathbb{N})$
- 7. State whether  $\hat{\mathcal{R}}(E \cup \{1\}) \subseteq E \cup \{1\}$

You may which to exploit the answer for some question when answering another. Finally:

- 1. Characterize the minimum fixed point of  $\hat{\mathcal{R}}$ , i.e.  $\bigcap \{X \mid \hat{\mathcal{R}}(X) = X\}$
- 2. Characterize the maximum fixed point of  $\hat{\mathcal{R}}$ , i.e.  $\bigcup \{X \mid \hat{\mathcal{R}}(X) = X\}$

### 4.1 Answer

Write your answer here.