

# Computability Assignment

## Year 2012/13 - Number 2

Please keep this file anonymous: do not write your name inside this file.

More information about assignments at <http://disi.unitn.it/~zunino/teaching/computability/assignments>

### 1 Question

In this exercise,  $p(x)$  and  $q(x)$  will be two unary properties over natural numbers, and  $P$  and  $Q$  will denote the sets  $P = \{x \in \mathbb{N} : p(x) \text{ holds}\}$  and  $Q = \{x \in \mathbb{N} : q(x) \text{ holds}\}$ . If possible, for each of the cases below find two properties  $p(x)$  and  $q(x)$  such that  $\forall x \in \mathbb{N}. p(x) \Rightarrow q(x)$  and

1.  $P \subset Q$  (strict inclusion);
2.  $Q \subset P$  (strict inclusion);
3.  $P \setminus Q \neq \emptyset$ ;
4.  $Q \setminus P \neq \emptyset$ .

If for some of the above cases it's impossible to find such properties, provide a brief explanation of why is it so.

#### 1.1 Answer

1.  $p(x) = (x \text{ is divisible by } 4)$  and  $q(x) = (x \text{ is even})$ . Then,  $\forall x \in \mathbb{N}. p(x) \Rightarrow q(x)$  and  $P \subset Q$  as  $P$  is always smaller than  $Q$ .
2. It is not possible.  $\forall x \in \mathbb{N}. p(x) \Rightarrow q(x)$  implies that for each element in  $P$ , it must also be in  $Q$ . So, either  $P$  is subset of  $Q$  or  $P$  is equal to  $Q$ .
3. It is not possible. From the argument as explained in 2,  $P$  is subset of  $Q$ . Hence,  $P \setminus Q$  is always empty.
4. The same example as 1. As set  $Q$  is always larger than  $P$ ,  $Q \setminus P \neq \emptyset$

## 2 Preliminaries

Given an infinite sequence of sets  $(A_i)_{i \in \mathbb{N}}$ , we define  $\bigcap_{i=0}^{\infty} A_i = \bigcap \{A_i \mid i \in \mathbb{N}\} = \{x \mid \forall i \in \mathbb{N} x \in A_i\}$  and  $\bigcap_{i=0}^k A_i = \bigcap \{A_i \mid i \in \mathbb{N} \wedge i \leq k\} = A_0 \cap A_1 \cap \dots \cap A_k$ .

## 3 Question

Assume  $(A_i)_{i \in \mathbb{N}}$  to be an infinite sequence of sets of natural numbers, satisfying

$$\mathbb{N} \supseteq A_0 \supseteq A_1 \supseteq A_2 \supseteq A_3 \cdots (*)$$

For each property  $p_i$  shown below, state whether

- the hypothesis  $(*)$  is sufficient to conclude that  $p_i$  holds; or
- the hypothesis  $(*)$  is sufficient to conclude that  $p_i$  does not hold; or
- the hypothesis  $(*)$  is not sufficient to conclude anything about the truth of  $p_i$ .

Justify your answers (briefly).

1.  $p_1$ :  $\forall k \in \mathbb{N}. A_k = \bigcap_{i=0}^k A_i$ ;
2.  $p_2$ : if  $\forall i \in \mathbb{N}. A_i$  is finite, then there exists  $j \in \mathbb{N}$  such that  $A_j = A_{j+1}$ ;
3.  $p_3$ : for all  $i$ , if  $A_i$  is finite, then  $A_i = A_{i+1}$ ;
4.  $p_4$ : if  $\forall i \in \mathbb{N}. A_i \neq A_{i+1}$ , then  $\bigcap_{i=0}^{\infty} A_i = \emptyset$ ;
5.  $p_5$ : if  $\forall i \in \mathbb{N}. A_i$  is finite, then  $\bigcap_{i=0}^{\infty} A_i$  is finite;
6.  $p_6$ : if  $\forall i \in \mathbb{N}. A_i$  is infinite, then  $\bigcap_{i=0}^{\infty} A_i$  is finite;
7.  $p_7$ : if  $\forall i \in \mathbb{N}. A_i$  is infinite, then  $\bigcap_{i=0}^{\infty} A_i$  is infinite.

### 3.1 Answer

1. Sufficient to conclude that  $p_i$  holds. As  $A_k$  is subset of all the sets  $A_0$  to  $A_{k-1}$ , all the elements of  $A_k$  should belong to all those sets. It can not have any extra element not belonging to all the sets before  $A_k$
2. Sufficient to conclude that  $p_i$  holds. Since all the sets  $A_i$  are finite, the possible unique subsets of the set is also finite. But there are infinite  $A_i$ , hence some of the  $A_i$  must repeat. Also, from the condition that  $A_{k-1} \supseteq A_k$ , only the consecutive sets could be equal. Hence this property is true.

3. Insufficient. If  $A_i = \{2, 3, 4, 5\}$ , then from the given condition we can say  $A_i \supseteq A_{i+1}$  which means  $A_{i+1}$  can be  $\{2, 3, 4, 5\}$  in which case  $A_i = A_{i+1}$  is satisfied. But,  $A_{i+1}$  can also be  $\{2, 3, 4\}$  in which case  $A_i = A_{i+1}$  is not true.
4. Insufficient.
5. Sufficient to conclude that  $p_i$  holds. The intersection of finite set is always finite.
6. Sufficient to conclude that  $p_i$  does not hold. As all sets are infinite and each  $i+1$  th set is subset of  $i$  th set, the intersection should be infinite.
7. Sufficient to conclude that  $p_i$  holds