

Computability Assignment

Year 2013/14 - Number 6

Please keep this file anonymous: do not write your name inside this file.

More information about assignments at <http://disi.unitn.it/~zunino/teaching/computability/assignments>

Please do not submit a file containing only the answers; edit this file, instead, filling the answer sections.

1 Question

Remember that for all $A \subseteq \mathbb{N}$, $\overline{A} = \mathbb{N} \setminus A$, and id_A is the identity function on A .

Let $f \in (\mathbb{N} \rightarrow \mathbb{N})$ and let $A = \{f(n) \mid n \text{ is a prime number}\}$.

1. Characterize the elements of the set \overline{A} (i.e. find a property p such that $\overline{A} = \{n \mid p(n)\}$). Notice that p could be a conjunction of many “simpler” properties.
2. Define a function $g \in (A \rightarrow \mathbb{N})$ such that $f \circ g = \text{id}_A$.

1.1 Answer

1. We define the property p as $p \equiv (n = f(m) \text{ for some } n \in \mathbb{N}) \wedge (m = pq \text{ such that } p, q \neq 1 \wedge p, q \in \mathbb{N})$.
2. $g(a) = \min(\{m \in \mathbb{N} \mid f(m) = a\})$. Note g is property defined for each element of A .

2 Question

Let $A = \{n \mid \exists m \in \mathbb{N}. n = m^2\}$ and $B = \{2n \mid n \in \mathbb{N}\}$. Following the steps outlined below, define a bijection $f \in (\mathbb{N} \rightarrow \mathbb{N})$ such that $f(A) = B$ and $f(\overline{A}) = \overline{B}$.

1. Provide a bijection $g \in (A \rightarrow \mathbb{N})$.
2. Provide a bijection $h \in (\mathbb{N} \rightarrow B)$.

3. Argue that there exists a bijection $g' \in (\overline{A} \rightarrow \mathbb{N})$.
4. Provide a bijection $h' \in (\mathbb{N} \rightarrow \overline{B})$.
5. Prove that the function $f \in (\mathbb{N} \rightarrow \mathbb{N})$ defined as

$$f(n) = \begin{cases} (h \circ g)(n) & \text{if } n \in A \\ (h' \circ g')(n) & \text{if } n \in \overline{A} \end{cases}$$

satisfies all the desired properties.

2.1 Answer

1. $g(a) = \sqrt{a}$ (note that, by definition of A , $\forall a \in A. \sqrt{a} \in \mathbb{N}$).
2. $h(a) = 2 \cdot a$.
3. It is always possible to find a bijection between \mathbb{N} and any non-finite $D \subseteq \mathbb{N}$. This can be done by enumerating.
4. $h'(n) = 2 \cdot n + 1$.
5. Consider the two cases for which f is defined:
 - (a) if $n \in A$, then $f(n) = (h \circ g)(n) = h(g(n)) = 2 \cdot \sqrt{n}$. Thus $f(A) = h(g(A)) = h(\mathbb{N}) = B$. Note this is a bijection (composition of two bijections);
 - (b) if $n \in \overline{A}$, then $f(n) = (h' \circ g')(n) = h'(g'(n)) = 2 \cdot g'(n) + 1$. Note that, by construction, $g'(\overline{A}) = \mathbb{N}$, so $f(\overline{A}) = h'(g'(\overline{A})) = h'(\mathbb{N}) = \overline{B}$; note this is a bijection as well.

Note f is the union of 2 bijections between disjoint sets, so it is itself a bijection.